

Matter Effects of Thin Layers: Detecting Oil by Oscillations of Solar Neutrinos

Ara N. Ioannisian^{1,3,4} and Alexei Yu. Smirnov^{1,2}

1) International Center for Theoretical Physics, 34100 Trieste, Italy

2) Institute for Nuclear Research, RAS, Moscow, Russia

3) Institute for Theoretical Physics and Modeling, Halabian 34, Yerevan-36, Armenia

4) Yerevan Physics Institute, Alikhanian Br. 2, Yerevan-36, Armenia

Abstract

We consider a possibility to use the solar neutrinos for studies of small scale structures of the Earth and for geological research. Effects of thin layers of matter with density contrast on oscillations of Beryllium neutrinos inside the Earth are studied. We find that change of the ${}^7\text{Be}$ neutrino flux can reach 0.1 % for layers with density of oil and size 20 km. Problems of detection are discussed. Hypothetical method would consist of measuring the ${}^7\text{Be}$ — flux by, *e.g.*, large deep underwater detector—submarine which could change its location.

1 Introduction

In geophysics the matter density profile of the Earth is determined by measurements of the seismic wave velocity profile. Seismic method is used widely also for the oil search, in particular, in the decision process before a trial well is drilled. This method has, however, shortcomings and alternatives are highly appreciated.

Enormous penetration ability of neutrinos is a source of strong temptation to use them in tomography and geological studies [1],[2],[3], [4]. The first proposal was based on inelastic scattering of high energy neutrinos produced by accelerators [1]. The authors have suggested several methods which use difference of the neutrino interactions in matter with different chemical composition and density.

Qualitatively different proposal is based on the elastic forward scattering (refraction) of neutrinos [2],[3],[4]. Refraction modifies properties of neutrino oscillations changing both the oscillation length and depth [5]. The effect depends on density of medium. Consequently, an appearance of layers with different densities on the way of neutrinos changes the oscillation pattern. In this connection, some possibilities to use the neutrino superbeams as well as the beams from neutrino factories have been considered [6].

There is a number of studies of the Earth matter effect on oscillations of solar neutrinos. It was marked that the solar neutrinos can give information about large scale density distribution inside the Earth. In particular, properties of the mantle and the core of the Earth can be studied [7].

The Earth matter effect has been considered for the ${}^7\text{Be}$ neutrinos [8] in connection to BOREXINO [10] and KamLAND [11] experiments. It was found that the effect is negligible for the LMA solution, at least for these detectors.

Main conditions to use neutrinos for the geological studies and searches for oil and minerals are:

- 1) sensitivity to small scale structures: $d \leq 100$ km;
- 2) possibility to move both a source and detector of neutrinos, so that substantial part of outer layers of the Earth mantle can be scanned.

No realistic proposals which satisfy these conditions have been published so far. In this paper we will consider a possibility which is based on:

- Detection of the solar ${}^7\text{Be}$ – neutrinos: For the LMA solution of the

solar neutrino problem the oscillation length of neutrinos with $E_{Be} \approx 0.86$ MeV is in the required range $l_\nu/2 \sim (10 \div 20)$ km.

- Measuring the time dependence of the neutrino flux by deep underwater detector - submarine.

The paper is organized as follows. In section 2 we consider the effect of thin layers of matter on neutrino oscillations (see [12] for some related work). We study dependence of the effect on neutrino parameters and properties of layers. In section 3 we consider a possibility to identify the effect of cavities. In section 4 we discuss problems of detections. Concluding remarks are given in section 5.

2 Effects of thin layers on neutrino oscillations

Let us consider a system of two mixed neutrinos with values of oscillation parameters Δm^2 and $\sin^2 2\theta$ from the LMA solution region [13]. (We will comment on the effect of the third neutrino at the end of this section). In the case of the LMA solution, the electron neutrino produced in the center of the sun is adiabatically converted to a combination of the mass eigenstates ν_1 and ν_2 which is determined by the mixing angle, θ_m^0 , in the production point:

$$\nu_e \rightarrow \cos \theta_m^0 \nu_1 + \sin \theta_m^0 \nu_2 . \quad (1)$$

The angle θ_m^0 is given by:

$$\tan^2 2\theta_m^0 = \tan^2 2\theta \left(1 - \frac{2V_e^0 E}{\Delta m^2 \cos 2\theta} \right)^{-2} , \quad (2)$$

where E is the neutrino energy, V_e^0 is the matter potential for neutrinos in the production point. $V_e = \sqrt{2}G_F\rho N_A Y_e$, G_F is the Fermi coupling constant, ρ is the density of medium, N_A is the Avogadro number and Y_e is the number of electrons per nucleon; $V_e^0 = V_e(\rho^0, Y_e^0)$.

Conversion effect should be averaged over the neutrino production region, and in what follows we will describe this averaging by the effective mixing angle $\bar{\theta}_m^0$.

Due to loss of coherence [14], neutrinos arrive at the surface of the Earth as incoherent fluxes of ν_1 and ν_2 with relative admixtures given by $\cos^2 \bar{\theta}_m^0$ and $\sin^2 \bar{\theta}_m^0$ correspondingly.

Let us consider oscillations inside the Earth. The probability to find ν_e in the detector can be written as:

$$P = \cos^2 \bar{\theta}_m^0 P_{1e} + \sin^2 \bar{\theta}_m^0 P_{2e} = \cos 2\bar{\theta}_m^0 P_{1e} + \sin^2 \bar{\theta}_m^0, \quad (3)$$

where P_{1e} and P_{2e} are the probabilities of $\nu_1 \rightarrow \nu_e$ and $\nu_2 \rightarrow \nu_e$ transitions in matter of the Earth correspondingly.

Suppose neutrinos cross the cavity with length d and density ρ_d which differs from the average density of surrounding matter, ρ . That is, the neutrinos cross consecutively three layers: with lengths and densities (l_1, ρ) , (d, ρ_d) , (l_2, ρ) . The total length of the trajectory is $L = l_1 + d + l_2$.

Let us find an effect of the cavity on oscillations. Introducing, P_{1e}^0 , the $\nu_1 \rightarrow \nu_e$ oscillation probability in absence of cavity (it corresponds to oscillations in the unique layer of density ρ and length L) we can write P as:

$$P = P^0 + \Delta P, \quad (4)$$

$$P^0 = \sin^2 \bar{\theta}_m^0 + \cos 2\bar{\theta}_m^0 P_{1e}^0, \quad (5)$$

$$\Delta P = \cos 2\bar{\theta}_m^0 (P_{1e} - P_{1e}^0). \quad (6)$$

Here P^0 is the probability to find ν_e in the absence of cavity and ΔP is the change of the probability due to the cavity effect.

We will calculate the effect for the beryllium neutrino line with energy $E_{Be} = 0.86$ MeV. For such an energy the matter effect on mixing and oscillations is very weak. It is determined by a small parameter:

$$\epsilon \equiv \frac{2V_e E}{\Delta m^2} \approx 2.6 \cdot 10^{-3} \left(\frac{\rho}{2.7 \text{ g/cm}^3} \right) \left(\frac{7 \cdot 10^{-5} \text{ eV}^2}{\Delta m^2} \right) \left(\frac{Y_e}{0.5} \right) \quad (7)$$

which characterizes deviations of the mixing angle and the oscillation length in medium from their vacuum values. We will use ϵ as an expansion parameter and find effects in the lowest order in ϵ . The oscillation length is determined by Δm^2 :

$$l_\nu^m \approx l_\nu \approx 30.7 \text{ km} \left(\frac{7 \cdot 10^{-5} \text{ eV}^2}{\Delta m^2} \right). \quad (8)$$

The probability of $\nu_1 \rightarrow \nu_e$ oscillations equals

$$P_{1e} = |\cos \theta S_{11} + \sin \theta S_{12}|^2, \quad (9)$$

where $S = ||S_{ij}||$ is the evolution matrix in the mass eigenstate basis, $\vec{\nu}$. The matrix S can be written as:

$$S = U D_2 U^\dagger U_d D_d U_d^\dagger U D_1 U^\dagger. \quad (10)$$

Here $U \equiv U(\rho)$ and $U_d \equiv U(\rho_d)$ are the mixing matrices of mass eigenstates in the layers 1, 2 and in the cavity correspondingly. They are defined in such a way that $\vec{\nu} = U \vec{\nu}_m$, where $\vec{\nu}_m$ is the basis of eigenstates of the Hamiltonian in matter. In (10), the diagonal matrices,

$$D_i \equiv \text{diag}(1, e^{i\Phi_i}), \quad i = 1, 2, d, \quad (11)$$

describe evolution of the neutrino eigenstates, $\vec{\nu}_m$, in layer i . Φ_i is the phase of oscillations:

$$\Phi_i = \frac{\Delta m^2 l_i}{2E} \sqrt{(\cos 2\theta - \epsilon_i)^2 + \sin^2 2\theta}, \quad (12)$$

$\epsilon_i \equiv \epsilon(\rho_i)$. In absence of cavity we have

$$S = S_0 = U D U^\dagger, \quad (13)$$

where $D = \text{diag}(1, e^{i\Phi})$ and

$$\Phi = \frac{\Delta m^2 L}{2E} \sqrt{(\cos 2\theta - \epsilon)^2 + \sin^2 2\theta} \quad (14)$$

is the total oscillation phase in absence of the cavity.

Using Eqs. (9,10,11), the straightforward calculation leads to the $\nu_1 \rightarrow \nu_e$ probability in the lowest order in ϵ :

$$P_{1e} \approx P_{1e}^0 + \epsilon \xi \sin^2 2\theta \sin \left(\Phi_2 + \frac{\Phi_d}{2} \right) \sin \frac{\Phi_d}{2}, \quad (15)$$

where

$$\xi \equiv \frac{(Y_e \rho) - (Y_{ed} \rho_d)}{Y_e \rho} \quad (16)$$

is the density contrast. In absence of cavity the probability equals

$$P_{1e}^0 = \cos^2 \theta - \epsilon \sin^2 2\theta \sin^2 \frac{\Phi}{2} . \quad (17)$$

And according to eq. (6) and (15) the effect of cavity can be written as

$$\Delta P = \epsilon \xi \cos 2\bar{\theta}_m^0 \sin^2 2\theta \sin \left(\Phi_2 + \frac{\Phi_d}{2} \right) \sin \frac{\Phi_d}{2} . \quad (18)$$

The following remarks are in order:

According to eq. (18) the effect of cavity is proportional to expansion parameter ϵ , the density contrast, ξ , and $\cos 2\bar{\theta}_m^0$. For maximal mixing in the production region the effect would be zero. For the best fit point of the LMA solution we get $\cos 2\bar{\theta}_m^0 \approx 0.3$.

Important feature of the result (18) is that ΔP does not depend on the oscillation effect in the first layer (before cavity), in particular, it does not depend on the phase Φ_1 . In contrast, ΔP does depend on Φ_2 – the oscillation phase in the layer between the cavity and detector. This property appears in the lowest order in ϵ and is related to the fact that initial state is the mass eigenstate and final state (at the detector) is the flavor state. It can be shown that if initial state is ν_e and final state – ν_1 , the probability ΔP does not depend on Φ_2 (phase between cavity and detector) but will depend on Φ_1 – the phase acquired in the first layer. Thus, the interchange of ν_1 and ν_e in the initial and final states leads to change $\Phi_2 \rightarrow \Phi_1$. This feature is important for identification of the cavity effect (see section 3).

Let us estimate the size of possible effect depending on parameters of neutrinos and the cavity. The cavity produces the change of the event rate during the time when the neutrino beam crosses the cavity. The relative value of the change $\Delta N/N = \Delta P/P$ equals in the lowest order in ϵ

$$\frac{\Delta P}{P^0} = f(\Delta m^2, \theta) \epsilon \xi \sin \left(\Phi_2 + \frac{\Phi_d}{2} \right) \sin \frac{\Phi_d}{2} , \quad (19)$$

where

$$f(\Delta m^2, \theta) = \frac{2 \cos 2\bar{\theta}_m^0 \sin^2 2\theta}{1 + \cos 2\bar{\theta}_m^0 \cos 2\theta} . \quad (20)$$

Here we have taken $P^0 = (1 + \cos 2\bar{\theta}_m^0 \cos 2\theta)/2$ neglecting the Earth matter effect. The dependence of $f(\Delta m^2, \theta)$ on $\tan^2 \theta$ for Δm^2 from the allowed

LMA region is shown in Fig. 1. According to Fig. 1 the function $f(\Delta m^2, \theta)$ has maximum $f^{max} \sim 0.4$ at about the best fit value of $\tan^2 \theta$.




Fig. 1. The dependence of $f(\Delta m^2, \theta)$ on $\tan^2 \theta$ for $\Delta m^2 = 5 \cdot 10^{-5} \text{ eV}^2$, $7 \cdot 10^{-5} \text{ eV}^2$ and 10^{-4} eV^2 .

The relative change of the probability can be rewritten as

$$\frac{\Delta P}{P^0} = A_{max} \sin \left(\Phi_2 + \frac{\Phi_d}{2} \right) \sin \frac{\Phi_d}{2}, \quad (21)$$

where

$$A_{max} \equiv \left. \frac{\Delta P_e}{P_e} \right|_{max} = \epsilon \xi f(\Delta m^2, \theta) \quad (22)$$

is the maximal value of $\Delta P/P^0$ which corresponds to both sines being 1. The effect increases with decrease of Δm^2 . For minimal allowed value $\Delta m^2 = 5 \cdot 10^{-5} \text{ eV}^2$, and $\tan^2 \theta = 0.3$, $\xi = 0.64$ we get

$$A_{max} = 0.1\%. \quad (23)$$

In Fig. 2 we show A_{max} as the function of Δm^2 and $\tan^2 \theta$. The effect weakly depends on θ in the allowed LMA region.




Fig. 2. Maximum value of the relative effect, $A_{max} \equiv (\Delta P/P)_{max}$, as the function of Δm^2 and $\tan^2 \theta$.

If detector is placed underwater, neutrinos will cross an additional layer of water, l_W , which can be rather large. Straightforward calculations lead to result, similar to eq.(19) in which

$$\Phi_2 \rightarrow \Phi'_2 = \Phi_2 + \Phi_w, \quad \Phi_w \simeq \frac{\Delta m^2 l_W}{2E},$$

where Φ_w is the phase acquired by neutrinos in water.

Let us comment on possible effect of the third neutrino. We consider the scheme in which ν_3 is separated from two others by the mass gap $\Delta m_{23}^2 = \Delta m_{atm}^2 = (2 - 3) \cdot 10^{-3} \text{ eV}^2$. (So, the scheme explains the atmospheric neutrino data.) We assume that this neutrino has small admixture of the electron neutrino described by $U_{e3} < 0.15$. The influence of matter on this mixing is determined by $\epsilon_{atm} = \epsilon(\Delta m_{atm}^2) \sim 7 \cdot 10^{-5}$. Consequently, variations of the flux due to cavity effect should be smaller than $7 \cdot 10^{-5}$. In fact, the observable effects are further suppressed by smallness of U_{e3} and averaging of oscillations associated with the third neutrino. For Δm_{atm}^2 the oscillation length is smaller than 1 km. Also interference between the modes of oscillations driven by Δm_{atm}^2 and solar Δm^2 produces negligible effect. So, we

conclude that the cavity effect is mainly due to oscillations driven by solar Δm^2 .

3 Effect of cavity. Exposure time

Let us consider the time dependence of the electron neutrino flux in, *e.g.*, deep underwater detector. For fixed position of the detector, the time dependence is due to rotation of the Earth. The rotation leads to certain change of the trajectory of solar neutrinos inside the Earth, and consequently, to modification of the oscillation effect. In general, the change of signal with time is both due to the cavity effect and the effect of regeneration in P^0 (see (5)). The problem is to disentangle these two effects.

Let t_{exp} be the exposure time - the time during which the trajectory of solar neutrinos crosses the cavity. The effect of cavity consists of an additional change of the neutrino flux during exposure time t_{exp} on the top of usual regeneration effect.

According to (5) and (17) the probability in the absence of cavity equals

$$P^0(t) = \sin^2 \bar{\theta}_m^0 + \cos 2\bar{\theta}_m^0 \left[\cos^2 \theta - \epsilon(t) \sin^2 2\theta \sin^2 \frac{\Phi(t)}{2} \right]. \quad (24)$$

In principle, this probability can be predicted with high enough accuracy. Indeed, the average radial density profile [15] is a subject to local variations at the level $(5 - 7)\%$ [16]. Therefore one expects $\sim (5 - 7)\%$ variations of $\epsilon(t)$. This uncertainty can be important for very small values of the density contrast: $\xi \sim (5 - 7)\%$ or very small phases in Eq. (19). Thus, one should look for the deviation of the time dependence of signal from the signal expected according to (24). Identification of the cavity effect depends on specific conditions of observation. If the cavity is close enough to the detector, the distance l_1 , and therefore L , can change significantly during the exposure time, so that the averaging over the phase Φ will occur. As a result

$$\overline{P^0} = \sin^2 \bar{\theta}_m^0 + \cos 2\bar{\theta}_m^0 \left(\cos^2 \theta - \frac{\epsilon}{2} \sin^2 2\theta \right) \quad (25)$$

practically does not depend on time, and whole time dependence is due to presence of cavity.

Due to smallness of relative change of the probability, $\Delta P/P$, one needs to collect the integral effect during long time of observation. Therefore, the effect of cavity is determined by the relative change of probability averaged over the exposure time:

$$\frac{\overline{\Delta P}}{P} = \frac{1}{t_{exp}} A_{max} \int dt \sin \left(\Phi'_2(t) + \frac{\Phi_d(t)}{2} \right) \sin \frac{\Phi_d(t)}{2} . \quad (26)$$

In general, the cavity has complicated form, so that $d = d(t)$ can change with time during exposure interval significantly. Averaging over d gives

$$\frac{\overline{\Delta P}}{P} = \frac{1}{t_{exp}} \frac{A_{max}}{2} \int dt \cos \Phi'_2(t) . \quad (27)$$

The size of the cavity should not be much smaller than the oscillation length. For the $\Delta m^2 \simeq 7 \cdot 10^{-5} \text{eV}^2$ the oscillation length is 30 km. Therefore maximum effect should be seen for the cavity size $l_\nu/2 \simeq 15$ km. For smaller cavities the effect will be suppressed according to the Eq. (19).

ΔP is the periodic function of the phase Φ'_2 . Averaging over Φ'_2 leads to disappearance of the cavity effect $\overline{\Delta P} = 0$. Suppose averaging is absent. By moving the detector one can vary distances l_2 and l_W . Selecting the distance l_2 and l_W in such a way that $\Phi'_2 + \Phi_d/2 = \pi/2 + \pi k$ ($k = \text{integer}$) one can maximize the effect

$$\Delta P = \epsilon \xi \sin^2 2\theta \cos 2\bar{\theta}_m^0 \sin \frac{\Phi_d}{2} . \quad (28)$$

Correspondingly, there are zones of l_2 with high and low sensitivity. Moving the detector by $l_\nu/2$ one can cover whole space.

The width of the beryllium neutrino line, $\Delta E_{Be} \approx 1.5 \text{ keV}$ [17], is very small. Consequently, no averaging over the energy occurs. Averaging would require the length of trajectory: $l_1 > l_\nu E_{Be}/\Delta E_{Be} \sim 2 \cdot 10^4 \text{ km}$ which is larger than the Earth diameter.

The sign of the effect depends on the sign of density contrast, the distance between the cavity and the detector, l_2 , and the size of the cavity itself. In particular, if the cavity is close enough to the detector, so that $\Phi_2 \ll \pi$, and therefore $\Delta P \sim \xi \sin^2 \Phi_d/2$, the ν_e flux (signal) increases, when neutrinos

cross the cavity for the positive contrast $\xi > 0$ $\rho_d < \rho_2$ (which is satisfied for oil or water), and it decreases for the negative contrast. Similar situation is for $\Phi_2 \sim 2\pi k$, $k = 1, 2, \dots$. The sign of the effect changes if the detector approaches or removes from the cavity by the distance $\sim l_\nu/2$.

Detecting the cavity from the first position of the detector one determines the line along which the cavity is situated. Observing the cavity from two different positions of detector one can reconstruct location of the cavity unless another cavity is detected from the second position. The latter can be checked by observation from the third position of the detector.

In general, one should take into account finite size of the region in which beryllium neutrinos are produced inside the sun. The diameter of the production region ($\sim 0.2R_\odot$) has the angular size at the Earth $\theta_{Be} = 9 \cdot 10^{-4}$ radian. Therefore, the cavity with transverse size h shields the production region completely if

$$l_2 \leq \frac{h}{\theta_{Be}} \simeq 1100 \text{ km} \left(\frac{h}{1 \text{ km}} \right). \quad (29)$$

If the cross-section of the cavity, S_d , is smaller than $(l_2\theta_{Be})^2$ an additional suppression factor appears in the size of the relative change of the flux:

$$\frac{\Delta P}{P^0} \propto \frac{S_d}{(l_2\theta_{Be})^2}. \quad (30)$$

The integral effect is determined by the *exposure time*, which in turn depends on (i) the size and shape of the cavity, (ii) its location, (iii) position of detector and (iv) season of the year. For simplicity we will consider the cavity of the parallelepiped form situated in the horizontal layer, with h being the thickness in the vertical plane, l_w — the width in the horizontal plane, and as before, d — the size of the cavity in horizontal plane along the direction from the Sun to the detector.

Let t_{exp}^{day} be the exposure time during the day, that is, during one crossing of the cavity by the neutrino trajectory. If ν — trajectory crosses the cavity N days during a year the total exposure time during the year equals:

$$t_{exp} \sim t_{exp}^{day} \cdot N. \quad (31)$$

Let us estimate the exposure time for different possible positions of the cavity.

Suppose the cavity is in the equatorial region. If the detector is situated in the same horizontal plane as the cavity, the diurnal exposure time is determined by the cavity thickness:

$$t_{exp}^{day} \sim \frac{t_{day}h}{2\pi l_2} \simeq 2.3 \min \left(\frac{h}{1\text{km}} \right) \left(\frac{100\text{km}}{l_2} \right), \quad (32)$$

where $t_{day} \equiv 24$ hours. If the detector is out of the plane or cavity is inclined, the exposure time can be larger. However, at the same time the length of ν trajectory in the cavity will be shorter. Both factors can compensate each other in the integral number events.

The cavity will shield the detector during $N = 365$ days, so that the total exposure time during the year can be as large as ~ 14 hours.

Let us consider the cavity near the North pole. Here the search will be possible in spring or in autumn when the sun appears at horizon. At this time the sun rises by $\sim 7 \cdot 10^{-3}$ radian/day. Therefore the sun will be shielded by the cavity during $N = h/(7 \cdot 10^{-3} l_2)$ days. For thickness $h = 1$ km we get $N = 1.5$ days. The diurnal exposure is determined by the width l_w , so that total exposure time equals

$$t_{exp} = \frac{l_w t_{day} N}{2\pi l_2} \text{day} \simeq 1 \text{ hour} \left(\frac{l_{width}}{20\text{km}} \right) \left(\frac{100\text{km}}{l_2} \right) \left(\frac{h}{1\text{km}} \right), \quad (33)$$

Let us consider now the search at latitudes 67° which correspond to the polar circle. In this case in December the sun is at horizontal (south) direction for 18 days a year. The diurnal exposure time is the same as in previous case. As a result, $t_{exp} = 13.5$ hours. Similar consideration holds for observation in June at the south latitude 67° .

4 Detection

Clearly, detection of so small ($< 0.1\%$) effect at low energies and during restricted time interval is a very challenging task. It may require new technological developments. Here we will give simple estimations which help to understand problems of the method.

Let us consider a possibility to detect the effect of the cavity by large underwater detector-submarine.

The background conditions should be as good as in the underground detectors of Beryllium neutrinos [10, 11]. To suppress the background generated by the cosmic rays, the detector should be placed deep underwater – at about 3500 m below sea level. This certainly restricts applications of the method and makes it more difficult technologically. Indeed, it is unlikely that, *e.g.*, oil reservoirs are very deep although such possibility is not excluded [18]. On the other hand, there is not too many places where the sea becomes very deep near the coast.

The background and therefore the depth could be reduced for detectors which use light elements (*e.g.*, helium) and efficient anti-coincidence shielding.

Let us estimate the change of number of events due to presence of cavity and required size of the detector. For definiteness we will consider detection of neutrinos via the νe^- scattering in scintillator. The rate of events due to the Beryllium neutrino flux without oscillations is [10]:

$$r_0 = 20 \frac{\text{events}}{\text{kt} \cdot \text{hour}} . \quad (34)$$

Then in presence of neutrino conversion the expecting rate is

$$r = \left[P + \frac{1}{5}(1 - P) \right] r_0 , \quad (35)$$

where P is the survival probability of the electron neutrinos. Here we have taken into account that $\nu_e e^-$ cross section is about 5 times larger than $\nu_\mu e^-$ cross section for ${}^7\text{Be}$ neutrino scattering with energy transfer (0.25 - 0.664) MeV. Taking typical value $P^0 \sim 0.6$ we get

$$r = 14 \frac{\text{events}}{\text{kt} \cdot \text{hour}} . \quad (36)$$

Total number of events detected during the year equals

$$N = r \cdot M_d \cdot t_{exp}^{year} , \quad (37)$$

where M_d is the mass of detector.

The effect of cavity consists of the increase (or decrease) of the number of events by amount

$$\Delta N_c = N \cdot \frac{\overline{\Delta P}}{P} \cdot \frac{1}{1 + 1/4P^0} = 0.7N \cdot \frac{\overline{\Delta P}}{P} \quad (38)$$

(see Eq. (26)). Last factor in this equation reflects dumping effect due to contribution of neutral current interactions. This factor is absent for detection via the charged current interactions.

Let us find the number of events needed to establish an excess (or deficit) of the number of event due to presence of the cavity at $k\sigma$ level (k is integer). We take an ideal situation of 100 % efficiency of detection, absence of background, absence of averaging, etc.. In this case the possibility to identify the effect of cavity will be determined by statistical fluctuation of the neutrino signal during the exposure time: $\Delta N_{stat} = \sqrt{N}$. The effect of cavity should be larger than the fluctuation. The effect of cavity at $k\sigma$ level will be achieved if $\Delta N_c \geq k\Delta N_{stat} = k\sqrt{N}$. From this condition we find using Eq. (38):

$$N \geq 2 \cdot k^2 \cdot \left(\frac{\Delta P}{P} \right)^{-2}. \quad (39)$$

For $k = 3$ (3σ - level) and $\Delta P/P = 0.001$ this inequality gives

$$N \geq 1.8 \cdot 10^7 \text{ events}. \quad (40)$$

Consequently, the mass of detector should be

$$M_d \geq \frac{N}{rt_{exp}} = \frac{2 \cdot k^2}{rt_{exp}} \cdot \left(\frac{\Delta P}{P} \right)^{-2}. \quad (41)$$

Taking the exposure time $t_{exp} \sim 10$ hours we find $M = 130$ Mt. In general,

$$M_d = 130 \text{ Mt} \frac{k^2}{9} \frac{10 \text{ hours}}{t_{exp}}. \quad (42)$$

For the density of scintillator 0.8 g/cc a detector could have dimensions $\sim (500 \text{ m})^3$. (Compare with SuperKamiokande which has diameter 39 m and the height 42 m.) Notice that for such a large detector the problem can appear with fiducial volume. In our estimations of the exposure time we have neglected the finite sizes of the detector. In fact, only part of the so large detector can be shielded by the cavity: that is, the trajectories of neutrinos which cross the cavity will cover only part of the detector during all exposure time. Therefore the effect of cavity will be further suppressed.

We have found that the size of detector (even in the most favorable situation) should be an order of magnitude larger than the size of future underground detectors which are now under consideration [19]. Solution of this problem may require next step of technological developments.

5 Discussion and conclusions

We have studied effects of the thin layers of matter on oscillations of solar neutrinos, in particular, on the Be-neutrinos. The effect is proportional to ϵ , the effective mixing parameter $\cos 2\bar{\theta}_m^0$ in the production point, and the density contrast.

The oscillations occur in the vacuum dominating regime and the matter effect appears as the small correction to the vacuum oscillations pattern. For the beryllium line the oscillation length is about 20 - 40 km and the matter effect parameter $\epsilon \sim 3 \cdot 10^{-3}$. This parameter put the absolute upper bound on the effect for a given energy. The only possibility to further enhance the effect is to have several layers with configurations which add up their effects coherently [20].

The interesting feature of the oscillation result is that for solar neutrinos it does not depend on the oscillation phase acquired by the neutrinos before the layer with density contrast.

We have considered a possibility to study small scale structures inside the Earth via oscillations of the solar Beryllium neutrinos. Such a study can be sensitive to cavities with density contrast $\sim O(1)$ and the size (10 - 20) km situated not too far ($<$ hundreds km) from the detector. The cavity can produce a change of the flux by about 0.1 % at most. We have found that exposure time during the year can reach about 10 hours. So, the detection of a cavity at 3σ level would require a detector of about 130 Mt size.

There are some ways to reduce the mass of detector:

- 1) The exposure time can be increased if the detector moves following the neutrino trajectories which cross the cavity. For this, of course, some *a priori* knowledge of possible position of the cavity is needed.

Another possibility is to perform observations in the same place during several years.

- 2) If sterile neutrinos participate in oscillations one can use coherent effects on scattering of neutrinos on nuclei by neutral currents in order to increase the cross section [21].

Refraction effect increases with neutrino energy: $\epsilon \propto E$. For the boron neutrinos we get $\epsilon \sim 0.03$. Moreover, the cross section of the νe -scattering is proportional to neutrino energy. Finally, the background conditions are better. So, one may think to use boron neutrinos instead of beryllium neutrinos. However, the flux of boron neutrinos is 3 orders of magnitude smaller, and therefore total number of events turns out to be one order of magnitude

smaller than in the case of beryllium neutrinos. There are other problems: the oscillation length is now about 300 km, and therefore with boron neutrinos one can probe larger structures than with $Be-$ neutrinos. Furthermore, due to continuous spectrum the problem appears with averaging over energies of neutrinos.

Further studies are needed to show feasibility of this method to study of small scale structures inside the Earth and for the geological research.

References

- [1] A. De Rujula, S. L. Glashow, R. R. Wilson and G. Charpak, Phys. Rept. **99**, 341 (1983).
- [2] V. K. Ermilova, V. A. Tsarev and V. A. Chechin, *SOV. PHYS. LEBEDEV INST. REP. (1988) NO. 3* 51-54. (*KRATKIE SOOB. FIZ. (1988) NO. 3* 38-40).
- [3] A. Nicolaidis, Phys. Lett. B **200**, 553 (1988).
- [4] A. Nicolaidis, M. Jannane and A. Tarantola, J. Geophys. Res. **96**, 21811 (1991).
- [5] L. Wolfenstein, Phys. Rev. D **17**, 2369 (1978). S. P. Mikheev and A. Y. Smirnov, Sov. J. Nucl. Phys. **42**, 913 (1985) [*Yad. Fiz.* **42**, 1441 (1985)]. S. P. Mikheev and A. Y. Smirnov, Nuovo Cim. C **9**, 17 (1986).
- [6] T. Ohlsson, W. Winter Phys. Lett. **B512** 357, (2001), e-Print Archive: hep-ph/0105293; T. Ohlsson, W. Winter, e-Print Archive: hep-ph/0111247.
- [7] see *e.g.*, E. Lisi, D. Montanino, Phys. Rev. **D56** 1792 (1997); J.N. Bahcall, P.I. Krastev, Phys. Rev. **C56** 2839 (1997); G.L. Fogli, E. Lisi, D. Montanino A. Palazzo, Phys. Rev. **D62** 113003 (2000); M. Maris and S. T. Petcov, arXiv:hep-ph/0004151; M. C. Gonzalez-Garcia, C. Pena-Garay and A. Y. Smirnov, Phys. Rev. D **63**, 113004 (2001).
- [8] A. de Gouvea, A. Friedland and H. Murayama, JHEP **0103**, 009 (2001) [arXiv:hep-ph/9910286].

- [9] V. D. Barger, D. Marfatia and B. P. Wood, Phys. Lett. B **498**, 53 (2001) [arXiv:hep-ph/0011251].
- [10] G. Ranucci *et al.* [BOREXINO Collaboration], Nucl. Phys. Proc. Suppl. **91**, 58 (2001).
- [11] "Proposal for US Participation in KamLAND", March 1999, <http://kamland.lbl.gov/KamLAND.US.Proposal.pdf>.
- [12] C. Lunardini, A.Yu. Smirnov, Nucl. Phys. **B583** 260 (2000); E. K. Akhmedov, Phys. Lett. **B503** 133 (2001); O. Yasuda, Phys.Lett. **B516** 111 (2001).
- [13] V. Barger, D. Marfatia, hep-ph/0212126; G.L. Fogli, E. Lisi, A. Marrone, D. Montanino, A. Palazzo, A.M. Rotunno, hep-ph/0212127; M. Maltoni, T. Schwetz, J.W.F. Valle, hep-ph/0212129; Abhijit Bandyopadhyay, Sandhya Choubey, Raj Gandhi, Srubabati Goswami, D.P. Roy, hep-ph/0212146; John N. Bahcall, M.C. Gonzalez-Garcia, Carlos Pena-Garay, hep-ph/0212147; H. Nunokawa, W.J.C. Teves, R. Zukanovich Funchal, hep-ph/0212202; P. de Holanda, A.Yu. Smirnov, hep-ph/0212270.
- [14] A. S. Dighe, Q. Y. Liu and A. Y. Smirnov, arXiv:hep-ph/9903329.
- [15] A. M. Dziewonski and D. L. Anderson, Phys. Earth Planet.Interiors **25**, 297 (1981).
- [16] R. J. Geller and T. Hara, arXiv:hep-ph/0111342.
- [17] A. V. Gruzinov, J. N. Bahcall, Astrophys. J. **490** 437 (1997).
- [18] Encyclopaedia Britannica, 15th ed., Chicago, University, 1982.
- [19] M. Aglietta, W. Fulgione, O. Saavedra, G. Trinchero, Nucl. Instrum. Meth. **A277** 17, (1989); Chang Kee Jun, hep-ex/000504.
- [20] V. K. Ermilova, V. A. Tsarev and V. A. Chechin, Kr. Soob. Fiz. *Short Notices of the Lebedev Institute*, **5** 26 (1986), E. Kh. Akhmedov, Yad. Fiz. **47** (1988) 475, P.I. Krastev, A.Yu. Smirnov, Phys. Lett. **B226** :341-346, 1989, P. I. Krastev, A. Yu. Smirnov, Mod. Phys. Lett. **A6**:1001, 1991.

- [21] A. Drukier and L. Stodolsky, Phys. Rev. **D 30**, 2295 (1984) L. Stodolsky, Phys. Rev. Lett. **34**, 110 (1975) [Erratum-ibid **34** 508 (1975)].